



THE IMPACT OF CAREGIVERS' INTERNAL WORKING MODELS ON CAREGIVING: A CASE STUDY ABOUT DISORDERS OF CONSCIOUSNESS

C. Romaniello^{1,2}, G. Grande¹, E. Bertoletti², V. Pedone^{2,3}, G. Northhoff⁴, M. Farinelli¹

¹ Clinical Psychology Service, Villa Bellombra Rehabilitation Hospital, Bologna, Italy. ² Santa Viola Hospital, Bologna, Italy. ³ Villa Bellombra Rehabilitation Hospital, Bologna, Italy. ⁴ Mind, Brain Imaging and Neuroethics, Institute of Mental Health Research, University of Ottawa, Ottawa, ON, Canada


INTRODUCTION AND BACKGROUND

-**Consciousness** can be defined by two components: arousal and awareness. Disorders of consciousness (DOCs) are characterized by a disrupted relationship between these two components.

-**Vegetative state**: defined by recovery of arousal without signs of awareness. It is characterized by:

- Wakefulness
- No awareness of self or environment
- No sustained, reproducible, purposeful behavioral responses to external stimuli
- No language comprehension or expression
- Relatively preserved hypothalamic and brain stem autonomic functions
- Bowel and bladder incontinence
- Variably preserved cranial-nerve and spinal reflexes (Gosseries et al., 2014)

Due to the severe compromising of autonomies, caregivers are crucial in both home and hospital settings.



In a vision in which the body (and the brain) perceives and simultaneously shapes environment and relationships, the caregiver may be considered as the relational context of the patient.

CAREGIVING AND ATTACHMENT

-Despite the severe impairment of consciousness, **the patient still represents a component of the attachment relationship** because he keeps on being present in caregivers' mind.

-**ATTACHMENT SYSTEM**: psychobiological system that motivates to build emotional bonds with significant others to protect themselves from threats and alleviate distress.

When the attachment figure is threatened (e.g. by an illness), caregiving behavior will be engaged to preserve the attachment bond.

-**Caregiving and attachment**: deeply inter-related (Bowlby, 1969; Simpson et al. 2010), both associated with specific neural circuitries (Panksepp, 1998; Coan, 2010). Caregiving behaviours are influenced by attachment styles (Romaniello et al., 2015) and Internal Working Models (Bowlby, 1969)

INTERNAL WORKING MODELS IN CAREGIVING

-Products of repetitive relational experiences from which individuals build unconscious mental representations and expectations about the self, significant others and the relationship between the two.

-Despite substantial changes DOCs cause, these caregiver's old intrapsychic and interpersonal dynamics keep on influence expectations, affective and behavioural interactions with the patient and the hospital professionals.

-To investigate the role of IWM and expectations about self and others in caregiving and to explore the relationships between IWM and other psychological features, a clinical case will be discussed.

CLINICAL CASE

-**Clinical Context**: a dedicated ward in S. Viola Hospital (Bologna, Italy), with a multidisciplinary staff, in which the caregiver is considered part of the care environment

SUBJECTS	
THE CAREGIVER	THE PATIENT
-Middle aged woman	-Middle aged man
-High school graduate	-High school graduate
-Secretary of her husband in a family-run business	-Medical salesman in a family-run business
-After the event, totally devoted to the care of her husband	-In a Vegetative State since 2007 after a heart attack
-Detached, alexithymic, neglected appearance, sceptical and wary with others	-Described as solitary, detached, usually far away from home
-Heavy smoker, Diabetes II type	-Severe hypertension
-Recent bereavement in the family (brother)	

REFERENCES

Beck AT et al. (1974) The measurement of pessimism: The hopelessness scale. *Journal of Consulting and Clinical Psychology*;42:861-865. **Beck, A.T et al. (1988)**. An inventory for measuring clinical anxiety: psychometric properties. *Journal of consulting and clinical psychology*, 56(6), 893. **Beck AT et al. (1996)**. Beck depression inventory-II. San Antonio. **Bond MP. (1995)**. The development and properties of the Defense Style Questionnaire. *Ego defenses: Theory and measurement*, 202-220. **Coan JA (2010)** Adult attachment and the brain. *Journal of Social and Personal Relationships*; 27:210-217. **Davis et al. (2003)**. The affective neuroscience personality scales: Normative data and implications. *Neuropsychology*, 5(1), 57-69. **Davis KL & Panksepp J. (2011)**. The brain's emotional foundations of human personality and the Affective Neuroscience Personality Scales. *Neurosci Biobehav Rev*, 35(9), 1946-1958. **Feeney et al. (1994)** Assessing adult attachment. In: Sperling MB, Berman WHM, editors. *Attachment in adults: Clinical and developmental perspectives*. New York: Guilford Press; pp 128-152. **Fossati A et al. (2003)** On the dimensionality of the Attachment Style Questionnaire in Italian clinical and nonclinical participants. *Journal of Social and Personal Relationships*; 20:55-79. **Ghisi M et al. (2006)**. Adattamento italiano di A.T. Beck et al. (1996). BDI-II, Beck Depression Inventory -II, Firenze: Organizzazioni Speciali. **Gosseries et al. (2014)**. Measuring consciousness in severely damaged brains. *Annual review of neuroscience*, 37, 457-478. **Marvardi M. et al. (2005)** Study Group on Brain Aging; Italian Society of Gerontology and Geriatrics. *Aging Clinical and Experimental Research*;17:46-53. **Novak M, Guest C.(1989)**. Application of a multidimensional caregiver burden inventory. *The Gerontologist*; 29:798-803. **Panksepp J** Affective Neuroscience: The foundations of human and animal emotions. New York: Oxford University Press; 1998. **Panksepp,J & Northhoff G. (2009)**. The trans-species core SELF: the emergence of active cultural and neuro-ecological agents through self-related processing within subcortical-cortical midline networks. *Consciousness and Cognition*, 18(1), 193-215. **Pompili M et al. (2009)** BHS Beck Hopelessness Scale: Manuale. Firenze: Giunti O.S. **Romaniello C., et al (2015)**. Anxious attachment style and hopelessness as predictors of burden in caregivers of patients with disorders of consciousness: A pilot study. *Brain injury*, 29(4), 466-472. **San Martini P. et al. (2004)**. Italian version of the defense style questionnaire. *Comprehensive psychiatry*, 45(6), 483-494. **Shiel et al., (2000)** The Wessex Head Injury Matrix (WHIM) main scale: a preliminary report on a scale to assess and monitor patient recovery after severe head injury. *Clinical Rehabilitation*, 14(4), 408-416. **Sica C. et al. (2005)**. Adattamento italiano di Beck, AT & Steer GK (1988). BAI, Beck Anxiety Inventory, Firenze, Giunti OS. **Sica C et al. (2008)**, Coping orientation of problem experienced - Nuova versione italiana (COPE-NVI): Uno strumento per la misura degli stili di coping. *Psicoterapia Cognitiva e Comportamentale*, 14(1), 27-53. **Simpson JA, Rholes WS (2010)**. Attachment and relationships: Milestones and future directions. *Journal of Social and Personal Relationships*, 27:173-180

INSTRUMENTS AND PROCEDURES

Beck Anxiety Inventory (BAI); Beck et al., 1988; Sica et al., 2005), a self-report measure of anxiety symptoms.

Semi-structured interview to collect caregiver and patient life history and features of their relationship

Affective Neuroscience Personality Scale (ANPS); Davis et al., 2003; Davis & Panksepp, 2011), a self-report measure of basic emotions (Panksepp, 1998): three positive (PLAY, SEEK, CARE), three negative (FEAR, ANGER, SADNESS).

Caregiver Burden Inventory (CBI); Novak & Guest, 1989; Marvardi et al.,2005) a self-report measure of burden



Defense Style Questionnaire (DSQ); Bond et al., 1995, San Martini et al., 2004), self-report measure of conscious derivatives of defense mechanisms grouped in immature and mature styles.

Beck Depression Inventory-II (BDI-2); Beck et al., 1996; Ghisi et al., 2006) a self-report measure of depressive symptoms.

Toronto Alexithymia Scale (TAS-20); Taylor et al., 1990, Bressi et al., 1996), a self-report measure of alexithymia through 3 scales: difficulty identifying and describing feelings, operational thinking.

Attachment Style questionnaire (Feeney et al., 1994; Fossati et al., 2003) a self-report measure of attachment through 5 dimensions: confidence, need for approval, discomfort with closeness, relationships as secondary, preoccupation with relationships

Coping orientations problems experienced (COPE); Sica et al., 2008), a self-report measure of behavioural strategies in stressful situations.



-Observation of interactions in daily life within the hospital setting

-Observation of structured multisensorial stimulation of the patient with self-related stimuli don by the caregiver

-Assessment of patient's basic interactions through the Wessex Head Injury Matrix (WHIM, Shiel et al., 2002; Di Stefano et al., 2012), a behavioural scale for monitoring active behaviours in DOCs.

RESULTS

SELF-REPORT	SCORES	AVERAGES/ CUT-OFF	RESULTS
BDI-II	31	13	Severe depression
BAI	17	13	Moderate Anxiety
COPE	32	M=23 SD=5	-High Avoiding
	19	M=27 SD=8.4	-Low social support
CBI	20	M=10,4 SD=5.4	-High time-dependent burden
TAS-20	65	61	-Alexithymia
ASQ	50	M=37 SD=7	-Avoidant Attachment
DSQ			-Immature defence profile (Splitting, Projection)
ANPS	33	M=28.25 SD=4.44	-Low SEEKING
	24	M=29.28 SD=4.44	-Low PLAY

OBSERVATION RESULTS

-Caregiver present in the hospital 9 hours per day every day, without vacations

-The interactions caregiver-patient is detached (third person speaking, patient called by his surname);

-Strenuous collaboration with hospital professionals

-Caregiver personally takes care of his hygiene and posture and she is the unique to deal with her husband hypertension


DISCUSSION

Caregiver shows a IWM typical of avoidant style attachment:

-**POSITIVE MODEL OF SELF** ("I will deal with threats alone") → high burden, difficult collaboration with hospital professionals

-**NEGATIVE MODEL OF OTHERS** ("Others are unreachable and unresponsive") → no seeking social support, low PLAY, DEACTIVATING STRATEGIES (avoiding coping strategy, alexithymia, external regulator of distress, psychosomatic pathologies).

BRAIN FUNCTIONING IN RELATIONSHIPS



A summary of the different levels of processing of the self in association with possible neural and psychological substrates in relation to different concepts of self. Also highlighted is the level of organization that may be shared between animals and humans (PROTO-SELF is preconscious) (Panksepp & Northhoff, 2009)